

タイトル	Iwanami Studies in Advanced Mathematics Hilbert Spaces of Analytic Functions Takahiko Nakazi Professor (Emeritus) of Hokkaido University Mathematical Subject Classification (2000): Primary 46E22; Secondary 47A15, 47B35, 60G10
著者	YAMAMOTO, Takanori
引用	北海学園大学学園論集(196): 125-134
発行日	2025-03-27

Iwanami Studies in Advanced Mathematics

Hilbert Spaces of Analytic Functions

Takahiko Nakazi

Professor (Emeritus) of Hokkaido University Mathematical Subject
Classification (2000): Primary 46E22; Secondary 47A15, 47B35, 60G10

Translated by Takanori YAMAMOTO

Introduction

This book is a commentary aimed at helping fourth-year undergraduate students in mathematics understand the space of holomorphic functions. A holomorphic function on the open unit disc in the complex plane can be expressed as a Taylor series there. Its radius of convergence is at least 1. The purpose of this book is to study a space where an inner product is defined in the linear space of such functions, and which becomes complete with respect to the norm determined by that inner product, namely a Hilbert space. Examples of Hilbert spaces of holomorphic functions include Hardy spaces, Bergman spaces, and Dirichlet spaces, all of which are named after people. These spaces are sometimes referred to by the same names even when they do not necessarily form Hilbert spaces, but the subject of this book is the case of Hilbert spaces. However, cases where they do not form Hilbert spaces are used as tools for that research. Here, the space of holomorphic functions is limited to the Hilbert space of holomorphic functions on the unit disc of one variable. Instead of the unit disc of a plane, we can consider more general planar domains or n -dimensional complex domains as the Hilbert space of holomorphic functions. However, the position of this book is to provide an easy-to-understand explanation of the Hilbert space of holomorphic functions on the unit disc of a plane, which is the simplest but has the most profound results. This book deals with a field that has made great progress from the first half of the 20th century to the present day in the 21st century, and continues to make progress. To go into a little more detail, a holomorphic function on an open unit disk in the complex plane can be expanded by Taylor expansion, and the radius of convergence is 1 or more. The purpose of this book is to study Hilbert spaces, which are linear spaces of such complex functions whose inner products are defined and which are complete with respect to the norm determined by these inner products. There have

been no books written to date that focus exclusively on Hilbert spaces, which are spaces of holomorphic functions. In that sense, this book is a new attempt. In the study of Hilbert spaces in this book, results on Banach spaces of holomorphic functions are often used. Since the subject of this book is Hilbert spaces in Part I, I would like first-time readers to finish Part I without going into the proofs of the theorems on Banach spaces in Part II that are cited. After that, I would like you to read the proofs in Part II for a deeper understanding. This book deals with Hilbert spaces with general reproducing kernels, which are very typical examples of Hilbert spaces of holomorphic functions and have a wide range of applications. Examples of such spaces include those defined by the coefficients of a Taylor expansion and those defined as the closure of polynomials. This book explains these two Hilbert spaces. Basic examples of Hilbert spaces of holomorphic functions are Hardy spaces, Bergman spaces, and Dirichlet spaces. These spaces have different functional analytical characteristics due to the different ways in which norms are implemented. Historically, the theory of Hardy spaces is the oldest, while the theories of Bergman spaces and Dirichlet spaces are relatively new. Hardy spaces have many applications outside of mathematics, but we will not go into them in this book. Let's take a closer look at the contents of this book. Part I discusses Hilbert spaces of holomorphic functions, which is the main subject of this book, while Part II discusses Banach spaces of holomorphic functions. However, Part II is not a generalization of Part I, but is limited to only the theorems used in Part I. In Chapter 1 of Part I, we state the basic theorems about general Hilbert spaces and bounded linear operators on them, but in a minimal way, in preparation for reading Chapters 2 to 9. However, we do not use the difficult spectral decomposition theorem, and we write it in a self-contained manner. In Chapter 2 of Part I, we investigate Hilbert spaces of holomorphic functions, those defined by sequences and those defined as polynomial closures, and describe their relationship. We also discuss the boundedness of shift operators and point functionals. However, this chapter is also written in a self-contained manner, mainly dealing with material that will help you understand Chapters 3 to 9. The work of Helson and Szegő in 1960 is the problem of drawing the spectrum of a stationary stochastic process whose angle between the past and future is non-zero, as well as the problem of drawing the weights when the analytic projection is bounded on a weighted square integrable Lebesgue space. Both of these are topics on weighted Hardy spaces. Chapter 7 deals with prediction problems of stochastic processes, and Chapter 8 discusses the boundedness of analytical projections.

I would like to express my sincere gratitude to Professor Yoshikazu Giga, a member of the editorial committee of this series, for giving me the opportunity to write this book. I would like to express my sincere gratitude to many people for their help in completing this book. In particular,

Takanori Yamamoto and Masahiro Yamada read my work carefully and provided many valuable comments. I would like to express my deepest gratitude to them. In particular, I would not have been able to complete this book without the assistance of Takanori Yamamoto. I would also like to thank Rikio Yoneda and Kohei Izuchi for listening to my seminars about the contents of this book. I would also like to thank the students at Hokkaido University who listened to the lecture and my wife, Keiko, for typing it up.

October, 2009

Takahiko Nakazi

Chapter 1 Hilbert space

This chapter describes the basic theorems for Hilbert space and its bounded linear operators, which are used in Chapters 2 and onwards of I. Proofs are given in all cases, except in unavoidable cases, but some use the theorem in Chapter 1 of II. With the exception of Section 8, the subject matter is classic, but Section 8 deals with two recent generalizations of isometric operators. This is used in Chapter 3.

Section 1 Definition of Hilbert space

Section 2 Orthogonal Projection

Section 3 Orthogonal Basis

Section 4 Bounded Linear Functional

Section 5 Bounded Linear Operator

Section 6 Compact Operator

Section 7 Square Root of Positive Linear Operator

Section 8 Operators Close to Isometric Operators

Section 1 Definition of Hilbert space

Definition I-1.1.1. $\mathcal{H}$ is a linear space with complex numbers $\mathbb{C}$ as scalars. For $f, g \in \mathcal{H}$, if $\langle f, g \rangle \in \mathbb{C}$, then $\langle \cdot, \cdot \rangle$ is called an inner product if it satisfies the following axioms.

- (1) $\langle f, f \rangle \geq 0$ ($f \in \mathcal{H}$), and the necessary and sufficient condition for $\langle f, f \rangle = 0$ is $f = 0$.
- (2) $\overline{\langle f, g \rangle} = \langle g, f \rangle$ ($f, g \in \mathcal{H}$), where $\overline{}$ denotes the complex conjugate.
- (3) $\langle f, g \rangle$ is linear in f and conjugate linear in g .

A linear space with an inner product defined is called an inner product space.

Lemma I-1.1.2. (Schwarz's inequality) Let $\langle \cdot, \cdot \rangle$ be the inner product on $\mathcal{H}$, and $\|f\| = \sqrt{\langle f, f \rangle}$ ($f \in \mathcal{H}$), then

$$|\langle f, g \rangle| \leq \|f\| \|g\| \quad (f, g \in \mathcal{H})$$

holds. Equality holds if and only if there exists $\alpha \in \mathbb{C}$ such that $g = \alpha f$ or $f = 0$.

Proof When $g = 0$, $\|g\| = 0$ and $\langle f, 0 \rangle = 0$, so both sides are 0 and the inequality holds. Therefore, let $g \neq 0$, and let $\alpha = \langle f, g \rangle / \|g\|^2$. Then

$$\begin{aligned} \|f - \alpha g\|^2 &= \langle f - \alpha g, f - \alpha g \rangle \\ &= \|f\|^2 - \alpha \langle g, f \rangle - \bar{\alpha} \langle f, g \rangle + \alpha \bar{\alpha} \|g\|^2 \\ &= \|f\|^2 - 2\operatorname{Re} \{ \alpha \langle g, f \rangle \} + |\alpha|^2 \|g\|^2 \\ &= \|f\|^2 - |\langle f, g \rangle|^2 / \|g\|^2 \end{aligned}$$

This completes the proof. ■

Theorem I-1.1.3. An inner product space $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ is a norm space with the norm defined as $\|f\| = \sqrt{\langle f, f \rangle}$. For the definition of the norm, see Definition II-1.1.1.

Proof It is sufficient to show that $\|\cdot\|$ is a norm. The axioms (1) and (2) of the norm are obvious. Now we show (3). According to Lemma I-1.1.2, for $f, g \in \mathcal{H}$, we have

$$\begin{aligned} \|f + g\|^2 &= \langle f + g, f + g \rangle \\ &= \|f\|^2 + 2\operatorname{Re} \langle f, g \rangle + \|g\|^2 \\ &\leq \|f\|^2 + 2\|f\| \|g\| + \|g\|^2 = (\|f\| + \|g\|)^2 \end{aligned}$$

This implies (3). This completes the proof. ■

Proposition I-1.1.4. Let $\mathcal{H}$ be an inner product space, and let $f, g \in \mathcal{H}$.

(1) (Midpoint Theorem)

$$\|f + g\|^2 + \|f - g\|^2 = 2(\|f\|^2 + \|g\|^2)$$

(2) (Polarization Identity)

$$\langle f, g \rangle = \frac{1}{4} (\|f + g\|^2 - \|f - g\|^2 + i\|f + ig\|^2 - i\|f - ig\|^2)$$

Proof Since

$$\|f \pm g\|^2 = \|f\|^2 \pm 2\operatorname{Re} \langle f, g \rangle + \|g\|^2,$$

(1) holds. Since

$$\|f + g\|^2 - \|f - g\|^2 = 4\operatorname{Re} \langle f, g \rangle,$$

$$\|f + ig\|^2 - \|f - ig\|^2 = 4\operatorname{Re} \langle f, ig \rangle = 4\operatorname{Im} \langle f, g \rangle,$$

(2) holds. This completes the proof. ■

Let $\|\cdot\|$ be a norm, and let $d(f, g) = \|f - g\|$. Then d is a distance, so a normed space is a metric space. For the distance, see Definition II-1.1.2.

Proposition I-1.1.5. In the inner product space, the inner product $\langle f, g \rangle$ is continuous with respect to f and g .

Proof Let $\|f_n - f_0\| \rightarrow 0$. Since $|\|f_n\| - \|f_0\|| \leq \|f_n - f_0\|$, we have $\|f_n\| \rightarrow \|f_0\|$, so $\{\|f_n\|\}$ is a bounded sequence. By the Schwarz inequality,

$$\begin{aligned} |\langle f_n, g_n \rangle - \langle f_0, g_0 \rangle| &\leq |\langle f_n, g_n - g_0 \rangle| + |\langle f_n - f_0, g_0 \rangle| \\ &\leq \|f_n\| \cdot \|g_n - g_0\| + \|f_n - f_0\| \cdot \|g_0\| \end{aligned}$$

Thus, if $\|f_n - f_0\| \rightarrow 0$ and $\|g_n - g_0\| \rightarrow 0$, then $\langle f_n, g_n \rangle \rightarrow \langle f_0, g_0 \rangle$ holds. ■

If $\lim_{n \rightarrow \infty} \|f_n - f\| = 0$, then it is called that f_n converges to f in norm. If $\lim_{n \rightarrow \infty} \langle f_n, g \rangle = \langle f, g \rangle$ for all $g \in \mathcal{H}$, then it is called that f_n weakly converges to f .

Proposition I-1.1.5 indicates that the norm convergent sequence is a weakly convergent sequence.

Definition I-1.1.6. An inner product space $\mathcal{H}$ is called a Hilbert space if it is complete with respect to the norm derived from its inner product.

Example I-1.1.7. Let X be a compact Hausdorff space. Let (X, Σ, μ) be the probability measure space, Let $L^2(X, \mu)$ be the set of all square-integrable functions with respect to μ on X ,

where two functions that coincide almost everywhere with respect to μ are identified. For $f, g \in L^2(X, \mu)$, let

$$\langle f, g \rangle = \int_X f \bar{g} d\mu.$$

Then $\langle f, g \rangle$ is an inner product on $L^2(X, \mu)$. $L^2(X, \mu)$ becomes a Hilbert space with respect to this inner product.

Proof By Theorem II-1.1.5, $L^2(X, \mu)$ is complete. Thus it is a Hilbert space. ■

Section 2 Orthogonal Projection

Let $\mathcal{H}$ be a Hilbert space. For $f, g \in \mathcal{H}$, let $\langle f, g \rangle = 0$. Then we say that f and g are orthogonal to each other, and we write $f \perp g$. A subset F, G of $\mathcal{H}$ is said to be orthogonal to each other if $\langle f, g \rangle = 0$ ($f \in F, g \in G$) holds, and we write $F \perp G$. Furthermore, the set of all elements that are orthogonal to F is denoted by $F^\perp$. $F^\perp = \{g \in \mathcal{H} : g \perp F\}$.

Definition I.-1.2.1. Let $\mathcal{H}$ be the inner product space, and let F be its closed subspace. For $h \in \mathcal{H}$, there is an $f \in F$ and a $g \in F^\perp$ such that $h = f + g$. Then f is called an orthogonal projection of h on F .

Theorem I.-1.2.2. Let $\mathcal{H}$ be a Hilbert space, and let F be its closed subspace. In this case, any $h \in \mathcal{H}$ can be uniquely decomposed into the form $h = f + g$ ($f \in F, g \in F^\perp$). That is, there is a unique orthogonal projection f on F of h .

Proof Uniqueness is clear. We show existence. Let

$$\delta = \text{dist}(h, F) = \inf_{l \in F} \|h - l\|.$$

Then, there exists $f_n \in F$ for $(n = 1, 2, \dots)$ such that $\|h - f_n\| \rightarrow \delta$ as $n \rightarrow \infty$. By the midpoint theorem I-1.1.4, since $(f_n + f_m)/2 \in F$, we have

$$\begin{aligned} 4\delta^2 + \|f_n - f_m\|^2 &\leq 4 \left\| h - \frac{f_n + f_m}{2} \right\|^2 + \|f_n - f_m\|^2 \\ &= \|(h - f_n) + (h - f_m)\|^2 + \|(h - f_n) - (h - f_m)\|^2 \\ &= 2\|h - f_n\|^2 + 2\|h - f_m\|^2. \end{aligned}$$

If $n \rightarrow \infty$ and $m \rightarrow \infty$, then $\|f_n - f_m\| \rightarrow 0$. Hence f_n is a Cauchy sequence. Since $\mathcal{H}$ is complete, there is an f such that $f = \lim_{n \rightarrow \infty} f_n \in \mathcal{H}$. Since F is closed, it follows that $f \in F$ and

$$\delta = \|h - f\| = \min_{l \in F} \|h - l\|.$$

Let $g = h - f$. We show that $g \perp F$. Let t be a real variable. For any $l \in F$,

$$X(t) = \|h - f - tl\|^2.$$

When $t=0$, the minimum value δ^2 is taken. Hence $X'(0)=0$. Since

$$X(t) = \|h - f\|^2 - 2t \operatorname{Re} \langle h - f, l \rangle + t^2 \|l\|^2,$$

it follows that $X'(0) = -2 \operatorname{Re} \langle h - f, l \rangle = 0$. We change l as il . Then $\langle h - f, l \rangle = 0 (l \in F)$. ■

Let $\mathcal{H}$ be a Hilbert space and let F be its closed subspace. Let $G = F^\perp$. Then G is also a closed subspace, and G is called an orthogonal complement of F . By Theorem I-1.2.2, we can show that $G^\perp = (F^\perp)^\perp = F$.

Section 3 Normal Orthogonal Basis

A countable subset $\{e_n\}$ of a Hilbert space $\mathcal{H}$ is called an orthonormal system of $\mathcal{H}$ if it satisfies the condition $\langle e_n, e_m \rangle = \delta_{nm}$, where δ_{nm} is the Kronecker delta. When the smallest closed subspace containing the orthonormal system $\{e_n\}$ denoted as

$$[\{e_n\}] = \{\text{closure of all linear combinations of } \{e_n\}\}$$

is equal to $\mathcal{H}$, then $\{e_n\}$ is called an orthonormal basis of $\mathcal{H}$.

Proposition I-1.3.1. Let $\{e_n\}$ be an orthonormal system of $\mathcal{H}$, and F be a closed subspace of $\mathcal{H}$ generated by $\{e_n\}$. For any $g \in \mathcal{H}$, if

$$f = \sum_{n=1}^{\infty} \langle g, e_n \rangle e_n,$$

then $f \in F$ and f is the orthogonal projection of g onto F .

References

1. V. M. Adamyan, D. Z. Arov and M. G. Krein, Infinite Hankel matrices and generalized Carathéodory-Fejer and Riesz problems, *Funct. Anal. Appl.* 2(1968), 1-18.

2. J. Agler, A disconjugacy theorem for Toeplitz operators, *Amer. J. Math.* 112(1990), 1–14.
3. A. Aleman, S. Richter and W. T. Ross, Pseudocontinuations and the backward shift, *Indiana Univ. Math. J.* 47(1998), 223–276.
4. E. Amar, Suites d'interpolation pour le classes de Bergman de la boule et du polydisque de $\mathbb{C}^n$, *Canad. J. Math.* 30(1978), 711–737.
5. C. Apostol, H. Bercovici, C. Foias and C. Pearcy, Invariant subspaces, dilation theory, and the structure of the predual of a dual algebra, I, *J. Funct. Anal.* 63(1985), 369–404.
6. S. Axler, The Bergman space, the Bloch space, and commutators of multiplication operators, *Duke Math. J.* 53(1986), 315–332.
7. N. Bari, "Biorthogonal systems and bases in Hilbert space", *Učeniye Zapiski, Moskovskii Ordena Lenina Gosudarslevennyi Universitet imeni M. V. Lomonosova*, Vol. 148, *Matematika*, 4(1951), pp. 69–107 (Russian); *Mathematical Reviews*, Vol. 14(1953), p. 289.
8. H. Bercovici, C. Foias, and C. Pearcy, Dual algebras with applications to invariant subspaces and dilation theory, *CBMS Regional Conference Series in Mathematics*, 56 ; published for the Conference Board of the Mathematical Sciences, Washington, D.C., by Amer. Math. Soc., Providence, R.I., 1985.
9. C. A. Berger, L. A. Coburn and K. H. Zhu, BMO on the Bergman spaces of the classical domains, *Bull. Amer. Math. Soc.* 17(1987), 133–136.
10. C. A. Berger, L. A. Coburn and K. H. Zhu, Function theory on Cartan domains and the Berezin-Toeplitz symbol calculus, *Amer. J. Math.* 110(1988), 921–953.
11. A. Beurling, On two problems concerning linear transformations in Hilbert space, *Acta Math.* 81 (1949), 239–255.
12. P. S. Bourdon, Cellular-indecomposable operators and Beurling's theorem, *Michigan Math. J.* 33(1986), 187–193.
13. L. Brown and A. L. Shields, Cyclic vectors in the Dirichlet space, *Trans. Amer. Math. Soc.* 285(1984), 269–304.
14. L. Carleson, An interpolation problem for bounded analytic functions, *Amer. J. Math.* 80(1958), 921–930.
15. L. Carleson, Interpolations by bounded analytic functions and the corona problem, *Ann. of Math.* 76 (1962), 547–559.
16. M. Cotlar and C. Sadosky, On the Helson-Szegő theorem and a related class of modified Toeplitz kernels, *Proc. Symp. Pure Math. Amer. Math. Soc.* 35(1979), 383–407.
17. C. Cowen and B. MacCluer, *Composition Operators on Spaces of Analytic Functions*, *Studies in Advanced Mathematics*, CRC Press, Boca Raton, FL, 1995.
18. A. Devinatz, Toeplitz operators on H^2 spaces, *Trans. Amer. Math. Soc.* 112(1964), 304–317.
19. J. Douglas, Solution of the problem of Plateau, *Trans. Amer. Math. Soc.* 33(1931), 263–321.
20. R. Douglas, H. Shapiro and A. Shields, Cyclic vectors and invariant subspaces for the backward shift operator, *Ann. Inst. Fourier (Grenoble)* 20(1970), 37–76.
21. P. L. Duren, *Theory of H^p Spaces*, Academic Press, New York., 1970.
22. P. Duren and A. Schuster, *Bergman Spaces*, American Mathematical Society, Providence, RI, 2004.
23. H. Dym and H. P. McKean, *Gaussian Processes, Function Theory, and the Inverse Spectral Problem*, Academic Press, New York., 1976.
24. T. W. Gamelin, *Uniform Algebras*, Prentice Hall, Englewood Cliffs, New Jersey, 1969.
25. J. B. Garnett, *Bounded Analytic Functions*, Academic Press, New York, 1981.
26. I. Gohberg, S. Goldberg and M. A. Kaashoek, *Classes of Linear Operators*, Vol. I, II T, *Operator Theory: Advances and Applications*, 49, Birkhäuser Verlag, Basel, 1990.

27. P. Hartman, On completely continuous Hankel matrices, *Proc. Amer. Math. Soc.*, 9(1958), 862–866.
28. W. W. Hastings, A Carleson measures theorem for Bergman spaces, *Proc. Amer. Math. Soc.* 52(1975), 237–241.
29. E. Hayashi, The kernel of a Toeplitz operator, *Integral Eq. Op. Th.* 9(1986), 588–591.
30. H. Hedenmalm, A factorization theorem for square area-integrable analytic functions, *J. Reine Angew. Math. J.* 4(1993), 163–174.
31. H. Hedenmalm, An invariant subspace of the Bergman space having the codimension two property, *J. Reine Angew. Math.* 443(1993), 1–9.
32. H. Hedenmalm, B. Korenblum and K. Zyu, *Theory of Bergman Spaces*, Springer-Verlag, New York, 2000.
33. H. Helson and D. Sarason, Past and future, *Math. Scand.* 21(1967), 5–16.
34. H. Helson and G. Szegő, A problem in prediction theory, *Annali di Mat. Pura et Applicata* 4, 51(1960), 107–138.
35. T. Hida, Canonical representations of Gaussian processes and their applications, *Mem. Coll. Sci. Univ. Kyoto* 33(1960), 109–155.
36. K. Hoffman, *Banach Spaces of Analytic Functions*, Prentice Hall, Englewood Cliffs, New Jersey, 1962.
37. P. Koosis, Moyennes quadratiques pondérées de fonctions périodiques et de leurs conjuguées harmoniques, *C. R. Acad. Sci. Paris* 291(1980), 255–257.
38. P. Koosis, *Introduction to H^p Spaces*, Cambridge Univ. Press, 1980.
39. D.H. Luecking, Trace ideal criteria for Toeplitz operators, *J. Funct. Anal.* 73(1987), 345–368.
40. G. McDonald and C. Sundberg, Toeplitz operators on the disk, *Indiana Univ. Math. J.* 28(1982), 229–237.
41. A. G. Miamee and M. Pourahmadi, Best approximations in $L^p(d\mu)$ and prediction problems of Szegő, Kolmogorov, Yaglom, and Nakazi, *J. London Math. Soc.* 38(1988), 133–145.
42. T. Nakazi, Two problems in prediction theory, *Studia Mathematica*, 78(1984), 7–14.
43. T. Nakazi, Kernels of Toeplitz operators, *J. Math. Soc. Japan* 38(1986), 607–616.
44. T. Nakazi, Bounded Hankel forms with weighted norms and lifting theorems, *Pacific J. Math.* 150 (1991), 123–137.
45. T. Nakazi, Toeplitz operators and weighted norm inequalities, *Acta Sci. Math. (Szeged)*, 58(1993), 443–452.
46. T. Nakazi, Isometric composition operators between two weighted Hardy spaces, *Nihonkai Math. J.* 17(2006), 111–124.
47. T. Nakazi and M. Yamada, Riesz’s function in weighted Hardy and Bergman spaces, *Canad. J. Math.* 48(1996), 930–945.
48. T. Nakazi and T. Yamamoto, Some singular integral operators and Helson-Szegő measures, *J. Funct. Anal.* 88(1990), 366–384.
49. T. Nakazi and T. Yamamoto, Weighted norm inequalities for some singular integral operators, *J. Funct. Anal.* 148(1997), 279–295.
50. T. Nakazi and T. Yamamoto, An argument of a function in $H^{1/2}$, preprint.
51. Z. Nehari, On bounded bilinear forms, *Ann. of Math.*, 65(1957), 153–162.
52. J. von Neumann, Allgemeine Eigenwerttheorie Hermitescher Funktionaloperatoren, *Math. Ann.* 102 (1929), 49–131.
53. J. Neuwirth and D. J. Newman, Positive $H^{1/2}$ functions are constant, *Proc. Amer. Math. Soc.* 18(1967), 958.
54. F. and R. Nevanlinna, Über die Eigenschaften analytischer Funktionen in der Umgebung einer

- singulären Stelle oder Linie, Acta Soc. Sci Fenn. 50(1922), No. 5.
55. M. Nishio, N. Suzuki and M. Yamada, Compact Toeplitz operators on parabolic Bergman spaces, preprint.
 56. V. L. Oleinik and B. S. Pavlov, Embedding theorems for weighted classes of harmonic and analytic functions, J. Soviet Math. 2(1974), 135–142.
 57. S. Richter, Invariant subspaces of the Dirichlet shift, J. Reine Angew. Math. 386 (1988), 205–220.
 58. S. Richter, A representation theorem for cyclic analytic two-isometries, Trans. Amer. Math. Soc. 328 (1991), 325–349.
 59. S. Richter and A. Shields, Bounded analytic functions in the Dirichlet space, Math. J. 198(1988), 151–159.
 60. S. Richter and C. Sundberg, A formula for the local Dirichlet integral, Michigan Math. J. 38(1991), 355–379.
 61. S. Richter and C. Sundberg, Invariant subspaces of the Dirichlet shift and pseudocontinuations, Trans. Amer. Math. Soc. 341(1994), 863–879.
 62. R. Rochberg, Toeplitz operators on weighted H^p spaces, Indiana Univ. Math. J., 26(1977), 291–298.
 63. R. Rochberg, Interpolation by functions in Bergman spaces, Mich. Math. J. 29(1982), 229–236.
 64. W. Rudin, Real and complex Analysis, Third Edition(1987), McGraw-Hill, Inc.
 65. D. E. Sarason, Generalized interpolation on H^∞ , Trans. Amer. Math. Soc. 127 (1967), 179–203.
 66. D. E. Sarason, An addendum to “Past and future”, Math. Scand. 30(1972), 62–64.
 67. K. Seip, Beurling type density theorems in the unit disk, Invent. Math. 113(1993), 21–39.
 68. K. Seip, Interpolation and Sampling in Spaces of Analytic Functions, University Lecture Series, Volume 33, Amer. Math. Soc., 2004.
 69. H. S. Shapiro and A. L. Shields, On some interpolation problems for analytic functions, Amer. J. Math. 83(1961), 513–532.
 70. A. L. Shields, Cyclic vectors in some spaces of analytic functions, Proc. Royal Irish. Acad. Sect. A 74 (1974), 293–296.
 71. S. M. Shimorin, Wold-type decomposition and wandering subspaces for operators closed to isometries, J. Reine Angew. Math., 531(2001), 147–189.
 72. G. Szegő, Beiträge zur Theorie der Toeplitzschen Formen, Math. Z. 6(1920), 167–202.
 73. T. Yamamoto, On the generalization of the theorem of Helson and Szegő, Hokkaido Math. J. 14(1985), 1–11.
 74. K. Zhu, Operator Theory in Function Spaces, Monographs and Textbooks in Pure and Appl. Math., 139, Marcel Dekker Inc, New York., 1990.
 75. K. Zhu, Evaluation operators on the Bergman space, Math. Proc. Comb. Phil. Soc. 117(1995), 513–523.